

## SM3H 2023-2024 Summer Packet

**Due Date: By first test in class. Approximately 2 weeks into 1<sup>st</sup> term. You will have in class assignments during this 2 week period as well.**

Hello students! We want to thank you for your interest in the secondary math 3 honors course. The function of this course is defined below:

Let  $x$  be a measure of student success with the SM2H curriculum and its prerequisites,

$$f(x) = \begin{cases} AP \text{ Calculus Student} & x = \text{Adequately prepared SM2H SM3H Master} \\ UVU 1010, \text{ Precalculus} & x = \text{Adequately prepared SM2 SM3 Master} \end{cases}$$

That is to say that the domain of this course includes students that have built a strong foundation with their SM2 experience and are prepared to move forward in mastering problem solving pertaining to trigonometry, polynomials with any real roots, conic sections, parametric equations, and the polar number system, to prepare to master the first year of calculus during the following year.

We've found that students often return to school in the fall with a layer of "rust" that needs to be scoured away in order to discern whether a student is compatible with an advanced math course at an honors pace. This packet's purpose is to help students show up on the first day without struggling to recover that which they have already mastered, while learning new curriculum.

This packet should contain **no new material**, provided you've had a strong learning experience in SM2. If you discover that your SM2 experience was lacking, you're welcome to contact us at [pstewart@alpinedistrict.org](mailto:pstewart@alpinedistrict.org) or [vshaw@alpinedistrict.org](mailto:vshaw@alpinedistrict.org) for assistance. If you are significantly confused by the material included in this packet, it might be time to reconsider your pursuit of the SM3H curriculum and look to SM3 or spending lots of time coming in for help.

Mr. Stewart & Miss. Shaw

OHS Math Dept

*I suggest using a 3-hole punch and inserting this packet into the same 3-ring binder that you'll use for note-taking and keeping materials for this course.*

## How to use this packet

We feel that the best results for students will come from following the pacing guide like the one shown below. Realize that this is just a **recommended** pacing; have fun over the summer! Adjust the pacing around your family vacations, movie marathons, or need to relax; just make sure your plan includes finishing the material by the **first test** of the 2021-2022 class.

You are looking to be a successful student AND have fun. Your SM2 experience should have covered the difference between the conjunctions AND and OR. Perhaps this summer will provide an opportunity to learn to manage your time to maximize both success AND fun. You'll certainly need time management skills to do well in your upper level academic course work.

### Week of summer "vacation"

Week of May 30  
Week of June 6  
Week of June 13  
Week of June 20  
Week of June 27  
Week of July 4  
Week of July 11  
Week of July 18  
Week of July 25  
Week of August 1  
Week of August 8

### Suggested Review Material


Pythagorean Theorem  
Trigonometric Functions  
  
Factoring Quadratic Expressions  
Factoring Special Cases  
Solve by Factoring  
  
Solve by Completing the Square  
Domain & Range

I suggest that plowing an entire worksheet in one sitting is not as impactful as sitting down for 20 minutes and doing 4 or 5 problems each day of the week. Having multiple exposures to a topic typically correlates well with assessments of how well the material "adheres" to the mind. Enjoy the light, leisurely pace; we won't **grade** this until we take the **first test**.

The following page contains a few helpful pointers for students that are struggling with recalling their lessons from SM2. Again, you are most welcome to contact us if you need a supplemental nudge in the right direction.

The packet will be graded by our first test and our first test will include some material from this packet. Consider this your first opportunity to be prepared for SM3

## Pythagorean Theorem




Named for the Greek mathematician, Pythagoras, this theorem relates the sides of a right triangle with the formula:

$$a^2 + b^2 = c^2$$

where  $a, b$  are the lengths of the legs of the right triangle and  $c$  is the length of the hypotenuse of the right triangle.


Points are represented by upper case letters. The side of the triangle opposite the angle at a given point is represented with the lower case version of the same letter (e.g., point B is directly across from the side with a length of  $b$ ).

Example 1: Find the missing side length of the triangle:



$$\begin{aligned} a^2 + b^2 &= c^2 && \text{Write the Pythagorean Theorem.} \\ 7^2 + 24^2 &= c^2 && \text{Use information from the figure.} \\ 49 + 576 &= c^2 && \text{Evaluate the exponentials.} \\ 625 &= c^2 && \text{Addition.} \\ \pm 25 &= c && \text{Square root each side.} \\ 25 &= c && \text{We prefer to consider distances as positive values, so we} \\ &&& \text{eliminate the negative solution.} \end{aligned}$$


Example 2: Find the missing side length of the triangle:



$$\begin{aligned} a^2 + b^2 &= c^2 && \text{Write the Pythagorean Theorem.} \\ a^2 + 9^2 &= 13^2 && \text{Use information from the figure.} \\ a^2 + 81 &= 169 && \text{Evaluate the exponentials.} \\ a^2 &= 88 && \text{Subtraction.} \\ a &= \pm\sqrt{88} && \text{Square root each side.} \\ a &= \pm 2\sqrt{22} && \text{Simplify the root.} \\ a &= 2\sqrt{22} && \text{We prefer to consider distances as positive values, so} \\ &&& \text{we eliminate the negative solution.} \end{aligned}$$

## Pythagorean Theorem

Find the missing side lengths of the right triangles using the Pythagorean Theorem given  $\triangle ABC$ :



1)  $a = 5, b = 6$ ; Find  $c$ .

2)  $a = 6, b = 8$ ; Find  $c$ .

3)  $a = 5, b = 12$ ; Find  $c$ .

4)  $a = 7, c = 25$ ; Find  $b$ .

5)  $a = 3, c = 12$ ; Find  $b$ .

6)  $a = 30, c = 34$ ; Find  $b$ .

7)  $b = 5, c = 5\sqrt{2}$ ; Find  $a$ .

8)  $b = 8, c = 16$ ; Find  $a$ .


9)  $b = 2.2, c = 8.7$ ; Find  $a$ .

## Trigonometric Functions

- A triangle with angle measurements of  $45^\circ$ ,  $45^\circ$ , and  $90^\circ$  has two legs of the same length and a hypotenuse with a length that is  $\sqrt{2}$  times the length of either leg.
- A triangle with angle measurements of  $30^\circ$ ,  $60^\circ$ , and  $90^\circ$  has two legs of different lengths. The longer of the legs has a leg that is  $\sqrt{3}$  times the length of the shorter leg. The length of the hypotenuse is twice as long as the shorter leg.
- A trigonometric ratio is a fraction that compares two sides of a triangle from the point of view of an angle. Each of the six trigonometric ratios has a unique comparison.
  - o Sine is the ratio of  $\frac{\text{Opposite Leg}}{\text{Hypotenuse}}$  and is written  $\sin \theta$ , where  $\theta$  is an angle.
  - o Cosine is the ratio of  $\frac{\text{Adjacent Leg}}{\text{Hypotenuse}}$  and is written  $\cos \theta$ , where  $\theta$  is an angle.
  - o Tangent is the ratio of  $\frac{\text{Opposite Leg}}{\text{Adjacent Leg}}$  and is written  $\tan \theta$ , where  $\theta$  is an angle.
  - o Cosecant is the ratio of  $\frac{\text{Hypotenuse}}{\text{Opposite Leg}}$  and is written  $\csc \theta$ , where  $\theta$  is an angle.
  - o Secant is the ratio of  $\frac{\text{Hypotenuse}}{\text{Adjacent Leg}}$  and is written  $\sec \theta$ , where  $\theta$  is an angle.
  - o Cotangent is the ratio of  $\frac{\text{Adjacent Leg}}{\text{Opposite Leg}}$  and is written  $\cot \theta$ , where  $\theta$  is an angle.

Example:

Find the value of each trigonometric ratio.



From  $\theta$ 's point of view:

- the opposite side's length is 15 units
- the adjacent side's length is 8 units
- the hypotenuse's length is 17 units



Therefore,

$$\sin \theta = \frac{15}{17}, \quad \cos \theta = \frac{8}{17}, \quad \tan \theta = \frac{15}{8}$$


$$\csc \theta = \frac{17}{15}, \quad \sec \theta = \frac{17}{8}, \quad \cot \theta = \frac{8}{15}$$

## Trigonometric Functions

Find the value of each segment or angle.



Find the value of each trigonometric ratio.



Find  $\sin \beta$


Find  $\cos \beta$

Find  $\tan \beta$

Find  $\csc \beta$

Find  $\sec \beta$

Find  $\cot \beta$



Find  $\sin \theta$

Find  $\cos \theta$

Find  $\tan \theta$

Find  $\csc \theta$

Find  $\sec \theta$

Find  $\cot \theta$

## Factoring Quadratic Expressions

- When factoring an expression of the form  $ax^2 + bx + c$ :
  - o if  $a = 1$ , then determine two factors (let's call them  $f_1$  and  $f_2$ ) of  $c$  that sum to  $b$  and write the factorization as:  $(x + f_1)(x + f_2)$ .
  - o if  $a \neq 1$ , then determine two factors (let's call them  $f_1$  and  $f_2$ ) of the product  $ac$  that sum to  $b$  and rewrite the expression as:  $ax^2 + f_1x + f_2x + c$ , then factor by grouping.

Example 1:

Factor  $x^2 + 5x - 24$

Since  $a = 1$ , I must find the factor pair of  $-24$  that adds up to  $+5$ .

The only factor pair of  $-24$  that sums to  $5$  is  $f_1 = 8, f_2 = -3$ .

So the factorization is  $(x + 8)(x - 3)$

Example 2:

Factor  $3x^2 + 11x + 10$

Since  $a \neq 1$ , I must find the factor pair of  $3 \cdot 10$  that adds up to  $+11$ .

The only factor pair of  $30$  that sums to  $11$  is  $f_1 = 5, f_2 = 6$ .

So we rewrite the original expression as  $3x^2 + 5x + 6x + 10$  and factor by grouping as follows:

$(3x^2 + 5x) + (6x + 10)$  Associate the first two terms and last two terms

$x(3x + 5) + 2(3x + 5)$  Factor the GCF from each association

$(x + 2)(3x + 5)$  Associate the "coefficients" of like terms

Your SM2 practice should have enabled you to casually glance at a quadratic expression and write the factorization down quickly. The above algorithms should only be executed if a student is struggling with a particular problem.

## Factoring Quadratic Expressions

Factor completely.

1)  $x^2 + 6x + 8$

2)  $x^2 - 7x - 18$

3)  $x^2 - 5x - 14$

4)  $x^2 - 16x + 63$

5)  $x^2 - 15x + 56$

6)  $x^2 - 9x + 8$

7)  $x^2 - 15x + 36$

8)  $x^2 + 22x + 40$

9)  $x^2 - x - 90$

10)  $2x^2 + 9x + 7$

11)  $3x^2 - 4x - 4$

12)  $5x^2 + 12x + 4$

13)  $4x^2 + 15x - 4$

14)  $10x^2 + 23x - 5$

15)  $6x^2 - 7x - 20$

16)  $30x^2 + 7x - 1$

17)  $12x^2 - 8x + 1$

18)  $4x^2 + 12x + 9$



## Factoring Special Cases

- When factoring an expression of the form  $a^2 - b^2$ :
  - Write the factorization as:  $(a + b)(a - b)$ .
- When factoring an expression of the form  $a^2 + 2ab + b^2$ :
  - write the factorization as:  $(a + b)(a + b)$  or  $(a + b)^2$
- When factoring an expression of the form  $a^2 - 2ab + b^2$ :
  - write the factorization as:  $(a - b)(a - b)$  or  $(a - b)^2$

Example 1:

Factor  $x^2 - 16$

As  $x^2$  and 16 are perfect squares and there is a - sign between them,  
the factorization is  $(x + 4)(x - 4)$

Example 2:

Factor  $4x^2 + 12x + 9$

As  $4x^2$  and 9 are perfect squares and the middle term is equal to twice the product of their roots ( $2 \cdot 2x \cdot 3 = 12x$ ),

the factorization is  $(2x + 3)^2$

Example 3:

Factor  $x^2 - 10x + 25$

As  $x^2$  and 25 are perfect squares and the middle term is equal to the opposite of twice the product of their roots ( $2 \cdot x \cdot (-5) = -10x$ ),

the factorization is  $(x - 5)^2$

For best results, remember to factor out the GCF of the expression before you do anything else!

## Factoring Special Cases

Factor completely.

1)  $x^2 - 16$

2)  $x^2 - 49$

3)  $x^2 - 121$

4)  $x^2 - 8x + 16$

5)  $x^2 - 10x + 25$

6)  $x^2 + 6x + 9$

7)  $x^2 - 100$

8)  $x^2 - 1$

9)  $x^2 - 400$

10)  $x^2 + 12x + 36$

11)  $x^2 - 7x + \frac{49}{4}$

12)  $x^2 + 20x + 100$

13)  $4x^2 - 64$

14)  $10x^2 - 90$

15)  $x^4 - 16$

16)  $4x^2 - 4x + 1$

17)  $100x^2 - 60x + 9$

18)  $4x^2 + 12x + 9$

## Solve by Factoring

- When solving a quadratic equation, bring all terms to one side of the equation so that the other side is zero. Then, factor the quadratic. As the product of multiple factors equaling zero is only possible by a factor being zero, set each factor equal to zero and solve these simpler equations to find the solutions to the original equation.

Example 1:

$$\text{Solve for } x: x^2 + 4x = -3$$

$$x^2 + 4x + 3 = 0 \quad \text{Bring all terms to the left side.}$$

$$(x + 3)(x + 1) = 0 \quad \text{Factor the quadratic.}$$

$$x + 3 = 0 \quad x + 1 = 0 \quad \text{Set each factor equal to zero.}$$

$$x = -3 \quad x = -1 \quad \text{Solve each equation.}$$

$$x = \{-3, -1\} \quad \text{Write the solutions to each equation in one solution set.}$$

Example 2:

$$\text{Solve for } x: x^3 + 4x^2 + x = 4x^2 + 26x$$

$$x^3 - 25x = 0 \quad \text{Bring all terms to the left side.}$$

$$x(x^2 - 25) = 0 \quad \text{Factor the GCF from the group.}$$

$$x(x - 5)(x + 5) = 0 \quad \text{Factor the quadratic.}$$

$$x = 0 \quad x - 5 = 0 \quad x + 5 = 0 \quad \text{Set each factor (including the GCF) equal to zero.}$$

$$x = 0 \quad x = 5 \quad x = -5 \quad \text{Solve each equation.}$$

$$x = \{-5, 0, 5\} \quad \text{Write the solutions to each equation in one solution set.}$$

## Solve by Factoring

Solve by factoring.

1)  $x^2 - 8x + 7 = 0$

2)  $x^2 + 6x + 5 = 0$

3)  $x^2 - 12x + 20 = 0$

4)  $6x^2 + 13x + 6 = 0$

5)  $x^2 - 36 = 0$



6)  $x^2 - 10x + 25 = 0$

7)  $2x^2 + 17x = -21$

8)  $7x^3 - 28x = 45x^2$


9)  $x^4 - 81 = 0$

10) Mrs. Shirk teaches in a rectangular classroom where the width of the classroom is 4 feet longer than the length of the classroom. The area of the classroom is 672 square feet. What are the dimensions of the classroom?



11) Mr. Elison has a rectangular garden that is 8 feet wide by 12 feet long. He decides to add a walkway around the garden that has the same width to form a larger rectangle. The area of the larger rectangle will be 224 square feet. What is the width of the walkway?


For each function or graph state the domain and range.



Domain:  
Range:

2)  $y = x^2 + 3x + 2$


Domain:  
Range:



Domain:  
Range:

4)  $y = 3x + 4$


Domain:  
Range:



Domain:  
Range:

6)  $y = |x - 4| + 3$


Domain:  
Range:



Domain:  
Range:

8)  $y = -2(x - 1)^2 + 3$

Domain:  
Range:



Domain:  
Range:

10)  $y = \begin{cases} x + 1, & x < -1 \\ 2, & x \geq 4 \end{cases}$

Domain:  
Range: